

## KUESIONER PENELITIAN

### 1. Identitas Responden

Mohon Saudara/i memberikan informasi demografi dengan mengisi titik-titik pada masing-masing pernyataan berikut ini:

1. Nama : .....
2. Usia : .....
3. Jenis Kelamin : .....
4. Masa kerja : .....
5. Pendidikan Terakhir : .....

### 2 Petunjuk Pengisian Point A

Karyawan dimohon untuk menjawab pernyataan-pernyataan di bawah ini dengan cara memberikan tanda *tick mark* / check (  $\surd$  ) pada salah satu jawaban untuk menyatakan persepsi. Setiap pernyataan dalam angket ini terdiri dari lima kategori jawaban yaitu :

- |        |                       |        |
|--------|-----------------------|--------|
| a. STS | = Sangat tidak setuju | Skor 1 |
| b. TS  | = Tidak Setuju        | Skor 2 |
| c. N   | = Netral              | Skor 3 |
| d. S   | = Setuju              | Skor 4 |
| e. SS  | = Sangat setuju       | Skor 5 |

### 3. Daftar Pernyataan :

#### a. Kinerja Karyawan

No	Pernyataan	Jawaban				
		STS	TS	N	S	SS
1	Karyawan mengerjakan semua pekerjaan dengan baik					
2	Karyawan mengerjakan pekerjaan dengan lengkap					
3	Karyawan membuat penjabaran tugas secara mandiri					
4	Karyawan mengerjakan tugas dengan penuh tanggung jawab					
5	Karyawan dapat melakukan penyesuaian diri dengan terhadap perubahan					
6	karyawan dapat bekerja sama dengan baik					

#### b. Gaya Kepemimpinan

No	Pernyataan	Jawaban				
		STS	TS	N	S	SS
	<b>Gaya Otoriter</b>					
1	Atasan memegang kekuasaan sendiri					
2	Kekuasaan terpusat pada atasan					
3	Rencana dirumuskan sendiri oleh atasan					
4	Kebijakan hanya dilakukan atasan					
5	Keputusan hanya dibuat atasan					
b.	<b><i>Lesuise faire</i></b>					
1	Atasan melimpahkan wewenang kepada bawahan					
2	Keputusan dibuat oleh bawahan					

3	Kebijakan dibuat oleh bawahan					
4	Komunikasi dilakukan sesuai keperluan					
5	Pengawasan terhadap bawahan agak longgar					
6	Pemikiran tentang pekerjaan datang dari atasan					
7	Pernan atasan sangat kecil dalam menyelesaikan pekerjaan					
8	Tanggung jawab dibebankan kepada masing-masing karyawan					
c.	<b>Demokratis</b>					
1.	Wewenang tidak mutlak dimiliki atasan					
2.	Wewenang dalam pekerjaan diberikan kepada masing-masing karyawan					
3.	Atasan menerapkan pendelegasian wewenang kepada bawahan					
4.	Ide dibuat secara musyawarah					
5.	Kebijakan dibuat berdasarkan hasil musyawarah					
6.	Kebijakan dibuat berdasarkan usulan bawahan					

## DATA PENELITIAN

## Data Kinerja (Y)

NO	Kode Responden	1	2	3	4	5	6	Jumlah
1	Kin1	4	4	3	4	5	5	25
2	Kin2	4	5	5	5	4	5	28
3	Kin3	4	4	4	4	5	4	25
4	Kin4	5	5	5	4	4	5	28
5	Kin5	5	4	4	4	3	4	24
6	Kin6	5	4	4	4	4	3	24
7	Kin7	4	4	5	4	5	4	26
8	Kin8	4	5	4	5	4	4	26

## Data Kepemimpinan Otoriter (X1)

NO	Kode Responden	1	2	3	4	5	Jumlah
1	Kin1	2	2	2	2	2	10
2	Kin2	3	2	3	2	1	11
3	Kin3	2	3	2	3	2	12
4	Kin4	3	3	3	2	3	14
5	Kin5	2	2	2	1	2	9
6	Kin6	2	2	2	2	2	10
7	Kin7	2	2	2	2	2	10
8	Kin8	1	3	3	2	2	11

Data Kepemimpinan *Laissez Faire* (X2)

NO	Kode Responden	1	2	3	4	5	6	7	8	Jumlah
1	Kin1	4	3	4	3	3	4	3	4	28
2	Kin2	3	4	4	4	4	3	4	4	30
3	Kin3	3	3	4	4	4	3	3	3	21
4	Kin4	4	4	3	3	3	4	4	3	21
5	Kin5	4	4	4	4	3	3	3	3	28
6	Kin6	4	4	3	3	4	4	3	4	29
7	Kin7	4	3	3	4	4	4	4	4	30
8	Kin8	4	3	4	4	4	4	4	4	31

Data Kepemimpinan Demokrasi (X3)

NO	Kode Responden	1	2	3	4	5	6	Jumlah
1	Kin1	4	4	3	4	5	5	25
2	Kin2	4	5	5	5	4	5	28
3	Kin3	5	4	4	4	5	4	26
4	Kin4	5	5	5	4	4	5	28
5	Kin5	5	4	3	4	3	4	23
6	Kin6	5	3	4	5	4	3	24
7	Kin7	4	4	5	4	5	4	26
8	Kin8	4	5	4	5	4	4	26

## HASIL UJI INSTRUMEN

### Reliability Statistics KINERJA

Cronbach's Alpha	Cronbach's Alpha Based on Standardized Items	N of Items
,909	,920	6

### Item-Total Statistics KINERJA

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Squared Multiple Correlation	Cronbach's Alpha if Item Deleted
kin1	21,0000	17,778	,747	,901	,897
kin2	20,7000	18,456	,703	,791	,903
kin3	20,0000	21,556	,812	,971	,894
kin4	20,2000	19,956	,771	,798	,891
kin5	20,4000	18,267	,841	,974	,879
kin6	20,2000	20,178	,740	,690	,895

### Reliability Statistics OTORITER

Cronbach's Alpha	Cronbach's Alpha Based on Standardized Items	N of Items
,892	,902	5

### Item-Total Statistics OTORITER

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Squared Multiple Correlation	Cronbach's Alpha if Item Deleted
go1	17,0000	8,667	,720	,582	,879
go2	16,8000	10,400	,847	,745	,858
go3	16,7000	8,900	,817	,738	,849
go4	16,8000	10,400	,653	,506	,886
go5	17,1000	9,433	,729	,566	,870

**Reliability Statistics Lesuise Faire**

Cronbach's Alpha	Cronbach's Alpha Based on Standardized Items	N of Items
,949	,950	8

**Item-Total Statistics Lesuise Faire**

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Squared Multiple Correlation	Cronbach's Alpha if Item Deleted
If1	29,1000	35,211	,877	,876	,937
If2	28,6000	36,267	,908	,987	,935
If3	28,5000	38,944	,741	,893	,946
If4	28,9000	38,544	,736	,928	,946
If5	28,7000	35,567	,902	,959	,935
If6	28,6000	34,711	,856	,968	,940
If7	28,6000	39,822	,821	,953	,943
If8	28,6000	40,933	,700	,857	,949

**Reliability Statistics DEMOKRATIS**

Cronbach's Alpha	Cronbach's Alpha Based on Standardized Items	N of Items
,887	,905	6

**Item-Total Statistics DEMOKRATIS**

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Squared Multiple Correlation	Cronbach's Alpha if Item Deleted
dk1	20,9000	12,989	,840	,949	,845
dk2	21,2000	13,067	,749	,797	,869
dk3	20,3000	18,900	,653	,726	,886
dk4	20,8000	16,622	,660	,848	,874
dk5	20,4000	16,933	,802	,941	,861
dk6	20,9000	15,656	,745	,748	,861

## HASIL ANALISIS DEKSRIPSI

### A. KINERJA

```
FREQUENCIES VARIABLES=Y1 Y2 Y3 Y4 Y5 Y6
  /NTILES=4
  /NTILES=10
  /STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN
  MEDIAN MODE SUM
  /ORDER=ANALYSIS.
```

### Frequencies

Notes	
Output Created	08-FEB-2019 22:20:19
Comments	
Input	Active Dataset DataSet0 Filter <none> Weight <none> Split File <none> N of Rows in Working Data File 8 Definition of Missing Missing Value Handling Cases Used
Syntax	User-defined missing values are treated as missing. Statistics are based on all cases with valid data. FREQUENCIES VARIABLES=Y1 Y2 Y3 Y4 Y5 Y6 /NTILES=4 /NTILES=10 /STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN MEDIAN MODE SUM /ORDER=ANALYSIS.
Resources	Processor Time 00:00:00,02 Elapsed Time 00:00:00,02

[DataSet0]



## Statistics

		Y1	Y2	Y3	Y4	Y5	Y6
N	Valid	8	8	8	8	8	8
	Missing	0	0	0	0	0	0
Mean		4,3750	4,3750	4,2500	4,2500	4,2500	4,2500
Std. Error of Mean		,18298	,18298	,25000	,16366	,25000	,25000
Median		4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
Mode		4,00	4,00	4,00	4,00	4,00	4,00
Std. Deviation		,51755	,51755	,70711	,46291	,70711	,70711
Variance		,268	,268	,500	,214	,500	,500
Range		1,00	1,00	2,00	1,00	2,00	2,00
Minimum		4,00	4,00	3,00	4,00	3,00	3,00
Maximum		5,00	5,00	5,00	5,00	5,00	5,00
Sum		35,00	35,00	34,00	34,00	34,00	34,00
10		4,0000	4,0000	3,0000	4,0000	3,0000	3,0000
20		4,0000	4,0000	3,8000	4,0000	3,8000	3,8000
25		4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
30		4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
40		4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
Percentiles	50	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
	60	4,4000	4,4000	4,4000	4,0000	4,4000	4,4000
	70	5,0000	5,0000	5,0000	4,3000	5,0000	5,0000
	75	5,0000	5,0000	5,0000	4,7500	5,0000	5,0000
	80	5,0000	5,0000	5,0000	5,0000	5,0000	5,0000
	90	.	.	.	.	.	.

## Frequency Table

Y1

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 4,00	5	62,5	62,5	62,5
Valid 5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

Y2

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 4,00	5	62,5	62,5	62,5
Valid 5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

Y3

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	1	12,5	12,5	12,5
Valid 4,00	4	50,0	50,0	62,5
Valid 5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

Y4

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 4,00	6	75,0	75,0	75,0
Valid 5,00	2	25,0	25,0	100,0
Total	8	100,0	100,0	

**Y5**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	1	12,5	12,5	12,5
4,00	4	50,0	50,0	62,5
5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

**Y6**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	1	12,5	12,5	12,5
4,00	4	50,0	50,0	62,5
5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

**B. KEPEMIMPINAN OTORITER**

```
FREQUENCIES VARIABLES=X1.1 X1.2 X1.3 X1.4 X1.5
  /NTILES=4
  /NTILES=10
  /STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN
MEDIAN MODE SUM
  /ORDER=ANALYSIS.
```

**Frequencies**

Notes		
Output Created		08-FEB-2019 22:40:10
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	8
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data.
Syntax		FREQUENCIES VARIABLES=X1.1 X1.2 X1.3 X1.4 X1.5
		/NTILES=4
		/NTILES=10
		/STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN MEDIAN MODE SUM
		/ORDER=ANALYSIS.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,03

[DataSet0]

## Statistics

		X1.1	X1.2	X1.3	X1.4	X1.5
N	Valid	8	8	8	8	8
	Missing	0	0	0	0	0
Mean		2,1250	2,3750	2,3750	2,0000	2,0000
Std. Error of Mean		,22658	,18298	,18298	,18898	,18898
Median		2,0000	2,0000	2,0000	2,0000	2,0000
Mode		2,00	2,00	2,00	2,00	2,00
Std. Deviation		,64087	,51755	,51755	,53452	,53452
Variance		,411	,268	,268	,286	,286
Range		2,00	1,00	1,00	2,00	2,00
Minimum		1,00	2,00	2,00	1,00	1,00
Maximum		3,00	3,00	3,00	3,00	3,00
Sum		17,00	19,00	19,00	16,00	16,00
Percentiles	10	1,0000	2,0000	2,0000	1,0000	1,0000
	20	1,8000	2,0000	2,0000	1,8000	1,8000
	25	2,0000	2,0000	2,0000	2,0000	2,0000
	30	2,0000	2,0000	2,0000	2,0000	2,0000
	40	2,0000	2,0000	2,0000	2,0000	2,0000
	50	2,0000	2,0000	2,0000	2,0000	2,0000
	60	2,0000	2,4000	2,4000	2,0000	2,0000
	70	2,3000	3,0000	3,0000	2,0000	2,0000
	75	2,7500	3,0000	3,0000	2,0000	2,0000
	80	3,0000	3,0000	3,0000	2,2000	2,2000
	90	.	.	.	.	.

## Frequency Table

### X1.1

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 1,00	1	12,5	12,5	12,5
2,00	5	62,5	62,5	75,0
3,00	2	25,0	25,0	100,0
Total	8	100,0	100,0	

### X1.2

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 2,00	5	62,5	62,5	62,5
3,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

### X1.3

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 2,00	5	62,5	62,5	62,5
3,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

### X1.4

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 1,00	1	12,5	12,5	12,5
2,00	6	75,0	75,0	87,5
3,00	1	12,5	12,5	100,0
Total	8	100,0	100,0	

**X1.5**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 1,00	1	12,5	12,5	12,5
2,00	6	75,0	75,0	87,5
3,00	1	12,5	12,5	100,0
Total	8	100,0	100,0	

### C. KEPEMIMPINAN LISUASE FAIRE

```

FREQUENCIES VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 X2.6 X2.7 X2.8
  /NTILES=4
  /NTILES=10
  /STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN
MEDIAN MODE SUM
  /ORDER=ANALYSIS.

```

### Frequencies

Notes		
Output Created		08-FEB-2019 22:35:01
Comments		
	Active Dataset	DataSet0
	Filter	<none>
Input	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	8
	Definition of Missing	User-defined missing values are treated as missing.
Missing Value Handling	Cases Used	Statistics are based on all cases with valid data.
		FREQUENCIES VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 X2.6 X2.7 X2.8 /NTILES=4 /NTILES=10 /STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN MEDIAN MODE SUM /ORDER=ANALYSIS.
Syntax		
	Processor Time	00:00:00,02
Resources	Elapsed Time	00:00:00,03

[DataSet0]



## Statistics

		X2.1	X2.2	X2.3	X2.4	X2.5	X2.6	X2.7	X2.8
N	Valid	8	8	8	8	8	8	8	8
	Missing	0	0	0	0	0	0	0	0
Mean		3,7500	3,5000	3,6250	3,6250	3,6250	3,6250	3,5000	3,6250
Std. Error of Mean		,16366	,18898	,18298	,18298	,18298	,18298	,18898	,18298
Median		4,0000	3,5000	4,0000	4,0000	4,0000	4,0000	3,5000	4,0000
Mode		4,00	3,00 <sup>a</sup>	4,00	4,00	4,00	4,00	3,00 <sup>a</sup>	4,00
Std. Deviation		,46291	,53452	,51755	,51755	,51755	,51755	,53452	,51755
Variance		,214	,286	,268	,268	,268	,268	,286	,268
Range		1,00	1,00	1,00	1,00	1,00	1,00	1,00	1,00
Minimum		3,00	3,00	3,00	3,00	3,00	3,00	3,00	3,00
Maximum		4,00	4,00	4,00	4,00	4,00	4,00	4,00	4,00
Sum		30,00	28,00	29,00	29,00	29,00	29,00	28,00	29,00
Percentile s	10	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000
	20	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000
	25	3,2500	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000
	30	3,7000	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000	3,0000
	40	4,0000	3,0000	3,6000	3,6000	3,6000	3,6000	3,0000	3,6000
	50	4,0000	3,5000	4,0000	4,0000	4,0000	4,0000	3,5000	4,0000
	60	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
	70	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
	75	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
	80	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
90	.	.	.	.	.	.	.	.	

a. Multiple modes exist. The smallest value is shown

## Frequency Table

**X2.1**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	2	25,0	25,0	25,0
Valid 4,00	6	75,0	75,0	100,0
Total	8	100,0	100,0	

**X2.2**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	4	50,0	50,0	50,0
Valid 4,00	4	50,0	50,0	100,0
Total	8	100,0	100,0	

**X2.3**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	3	37,5	37,5	37,5
Valid 4,00	5	62,5	62,5	100,0
Total	8	100,0	100,0	

**X2.4**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	3	37,5	37,5	37,5
Valid 4,00	5	62,5	62,5	100,0
Total	8	100,0	100,0	

**X2.5**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	3	37,5	37,5	37,5
Valid 4,00	5	62,5	62,5	100,0
Total	8	100,0	100,0	

**X2.6**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	3	37,5	37,5	37,5
Valid 4,00	5	62,5	62,5	100,0
Total	8	100,0	100,0	

**X2.7**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	4	50,0	50,0	50,0
Valid 4,00	4	50,0	50,0	100,0
Total	8	100,0	100,0	

**X2.8**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	3	37,5	37,5	37,5
Valid 4,00	5	62,5	62,5	100,0
Total	8	100,0	100,0	

## D. KEPEMIMPINAN DEMOKRATIS

```

FREQUENCIES VARIABLES=X3.1 X3.2 X3.3 X3.4 X3.5 X3.6
  /NTILES=4
  /NTILES=10
  /STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN
MEDIAN MODE SUM
  /ORDER=ANALYSIS.

```

## Frequencies

Notes		
Output Created		08-FEB-2019 22:26:02
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	8
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data.
Syntax		FREQUENCIES VARIABLES=X3.1 X3.2 X3.3 X3.4 X3.5 X3.6 /NTILES=4 /NTILES=10 /STATISTICS=STDDEV VARIANCE RANGE MINIMUM MAXIMUM SEMEAN MEAN MEDIAN MODE SUM /ORDER=ANALYSIS.
Resources	Processor Time	00:00:00,05
	Elapsed Time	00:00:00,03

[DataSet0]

## Statistics

		X3.1	X3.2	X3.3	X3.4	X3.5	X3.6
N	Valid	8	8	8	8	8	8
	Missing	0	0	0	0	0	0
Mean		4,5000	4,2500	4,1250	4,3750	4,2500	4,2500
Std. Error of Mean		,18898	,25000	,29505	,18298	,25000	,25000
Median		4,5000	4,0000	4,0000	4,0000	4,0000	4,0000
Mode		4,00 <sup>a</sup>	4,00	4,00 <sup>a</sup>	4,00	4,00	4,00
Std. Deviation		,53452	,70711	,83452	,51755	,70711	,70711
Variance		,286	,500	,696	,268	,500	,500
Range		1,00	2,00	2,00	1,00	2,00	2,00
Minimum		4,00	3,00	3,00	4,00	3,00	3,00
Maximum		5,00	5,00	5,00	5,00	5,00	5,00
Sum		36,00	34,00	33,00	35,00	34,00	34,00
Percentiles	10	4,0000	3,0000	3,0000	4,0000	3,0000	3,0000
	20	4,0000	3,8000	3,0000	4,0000	3,8000	3,8000
	25	4,0000	4,0000	3,2500	4,0000	4,0000	4,0000
	30	4,0000	4,0000	3,7000	4,0000	4,0000	4,0000
	40	4,0000	4,0000	4,0000	4,0000	4,0000	4,0000
	50	4,5000	4,0000	4,0000	4,0000	4,0000	4,0000
	60	5,0000	4,4000	4,4000	4,4000	4,4000	4,4000
	70	5,0000	5,0000	5,0000	5,0000	5,0000	5,0000
	75	5,0000	5,0000	5,0000	5,0000	5,0000	5,0000
	80	5,0000	5,0000	5,0000	5,0000	5,0000	5,0000
	90	.	.	.	.	.	.

a. Multiple modes exist. The smallest value is shown

## Frequency Table

### X3.1

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 4,00	4	50,0	50,0	50,0
Valid 5,00	4	50,0	50,0	100,0
Total	8	100,0	100,0	

### X3.2

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	1	12,5	12,5	12,5
Valid 4,00	4	50,0	50,0	62,5
Valid 5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

### X3.3

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	2	25,0	25,0	25,0
Valid 4,00	3	37,5	37,5	62,5
Valid 5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

### X3.4

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 4,00	5	62,5	62,5	62,5
Valid 5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

**X3.5**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	1	12,5	12,5	12,5
4,00	4	50,0	50,0	62,5
5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

**X3.6**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 3,00	1	12,5	12,5	12,5
4,00	4	50,0	50,0	62,5
5,00	3	37,5	37,5	100,0
Total	8	100,0	100,0	

## HASIL ANALISIS REGRESI

### Descriptive Statistics

	Mean	Std. Deviation	N
y	25,7500	1,58114	8
x1	10,8750	1,55265	8
x2	27,2500	3,99106	8
x3	25,7500	1,75255	8

### Correlations

		y	x1	x2	x3
Pearson Correlation	y	1,000	,684	-,147	,954
	x1	,684	1,000	-,732	,774
	x2	-,147	-,732	1,000	-,276
	x3	,954	,774	-,276	1,000
Sig. (1-tailed)	y	.	,031	,364	,000
	x1	,031	.	,019	,012
	x2	,364	,019	.	,254
	x3	,000	,012	,254	.
N	y	8	8	8	8
	x1	8	8	8	8
	x2	8	8	8	8
	x3	8	8	8	8

### Variables Entered/Removed<sup>a</sup>

Model	Variables Entered	Variables Removed	Method
1	x3, x2, x1 <sup>b</sup>	.	Enter

a. Dependent Variable: y

b. All requested variables entered.

### Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,962 <sup>a</sup>	,925	,869	,57244

a. Predictors: (Constant), X3, X2, X1



ANOVA<sup>a</sup>

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	16,189	3	5,396	16,468	,010 <sup>b</sup>
	Residual	1,311	4	,328		
	Total	17,500	7			

a. Dependent Variable: Y

b. Predictors: (Constant), X3, X2, X1

Coefficients<sup>a</sup>

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.	95,0% Confidence Interval for B		Correlations			Collinearity Statistics		
		B	Std. Error	Beta			Lower Bound	Upper Bound	Zero-order	Partial	Part	Tolerance	VIF	
		1	(Con)	1,182			4,174		,283	,791	-10,407	12,772		
1	x1	,095	,421	,093	,225	,833	-1,075	1,264	,684	,112	,031	,109	9,137	
1	x2	,070	,108	,178	,652	,550	-,229	,370	-,147	,310	,089	,253	3,959	
1	x3	,840	,265	,931	3,174	,034	,105	1,574	,954	,846	,434	,218	4,591	

a. Dependent Variable: y

Coefficient Correlations<sup>a</sup>

Model		x3	x2	x1	
1	Correlations	x3	1,000	-,675	-,874
		x2	-,675	1,000	,852
		x1	-,874	,852	1,000
	Covariances	x3	,070	-,019	-,097
		x2	-,019	,012	,039
		x1	-,097	,039	,177

a. Dependent Variable: y

**Collinearity Diagnostics<sup>a</sup>**


Model	Dimension	Eigenvalue	Condition Index	Variance Proportions			
				(Constant)	x1	x2	x3
1	1	3,965	1,000	,00	,00	,00	,00
	2	,032	11,104	,00	,02	,08	,00
	3	,002	47,548	1,00	,07	,29	,04
	4	,001	82,042	,00	,91	,63	,96

a. Dependent Variable: y


**Residuals Statistics<sup>a</sup>**


	Minimum	Maximum	Mean	Std. Deviation	N
Predicted Value	23,3161	27,8445	25,7500	1,52077	8
Std. Predicted Value	-1,600	1,377	,000	1,000	8
Standard Error of Predicted Value	,256	,491	,398	,082	8
Adjusted Predicted Value	22,6100	27,5295	25,6294	1,52620	8
Residual	-,62686	,68388	,00000	,43272	8
Std. Residual	-1,095	1,195	,000	,756	8
Stud. Residual	-1,660	1,719	,065	1,185	8
Deleted Residual	-1,44114	1,91864	,12065	1,10167	8
Stud. Deleted Residual	-2,579	2,909	,250	1,836	8
Mahal. Distance	,529	4,285	2,625	1,314	8
Cook's Distance	,002	2,070	,513	,717	8
Centered Leverage Value	,076	,612	,375	,188	8

a. Dependent Variable: y



Normal P-P Plot of Regression Standardized Residual







Mengambil + dari bapak Manager Koperasi Syariah Ar-Rahmah Banjarmasin



Menyerahkan Kuesioner kepada Kasir Koperasi Syariah Ar-Rahmah Banjarmasin

### RIWAYAT HIDUP

1. Nama : Anita
2. Tempat Tanggal Lahir : Negara , 07 Oktober 1995
3. Agama : Islam
4. Kebangsaan : Indonesia
5. Status Perkawinan : Belum Kawin
6. Alamat : Komplek Madya jaya Raya, Sungai Lulut .
7. Pendidikan :
  - a. SDN Bayanan 1 , Hulu Sungai Selatan Tahun 2008.
  - b. MTSN Tambak Bitin , Hulu Sungai Selatan Tahun 2011.
  - c. MAN Negara , Hulu Sungai Selatan Tahun 2014.
8. Pengalaman Organisasi : -
9. Orang Tua
  - Ayah :
  - Nama : Isra
  - Pekerjaan : Swasta
  - Alamat : Jl. Satria Tumbukan Banyu, Daha Selatan
  - Ibu
  - Nama : Syafiah (alm)
  - Pekerjaan : IRT
  - Alamat : jl. Satria tumbukan Banyu, Daha selatan
10. Anak Ke : 1 dari 4 Bersaudara

Banjarmasin, Juli 2019

Penulis

Anita